

FOURIER INVARIANT PARTIALLY APPROXIMATING SUBALGEBRAS OF THE IRRATIONAL ROTATION C*-ALGEBRA

S. WALTERS

The University of Northern British Columbia

May 30, 2001

ABSTRACT. For all transcendental parameters, the irrational rotation algebra is shown to contain infinitely many C*-subalgebras satisfying the following properties. Each subalgebra is isomorphic to a direct sum of two matrix algebras of the same (perfect square) dimension; the Fourier transform maps each summand onto the other; the corresponding unit projection is approximately central; the compressions of the canonical generators of the irrational rotation algebra are approximately contained in the subalgebra.

§1. Introduction

For each irrational number θ , the irrational rotation C*-algebra A_θ is generated by unitaries U, V satisfying the Heisenberg commutation relation $VU = \lambda UV$, where $\lambda = e^{2\pi i\theta}$. The Fourier transform on A_θ is the order four automorphism σ (or its inverse) of A_θ satisfying $\sigma(U) = V$ and $\sigma(V) = U^{-1}$. It can be viewed as a non commutative analogue of the Fourier transform of classical analysis (which itself plays an important role in the theory). Its square is the so-called flip automorphism and is fairly well understood (see [1], [3], [4], [5], [8], [14]). An open problem of Elliott (private communication), is if the Fourier transform has an inductive limit structure. More precisely, can A_θ be approximated by Fourier invariant basic building blocks consisting of finite direct sums of circle algebras and points? If it can be shown that the answer is affirmative, then it would seem that one is only a small step away from proving that the fixed point subalgebra A_θ^σ is an AF-algebra. One of the main difficulties in this problem (aside from some number theoretic ones), and which separates it from the corresponding flip problem, is that the projections needed in the construction are somewhat more exotic, as they can no longer be Rieffel projections. Therefore, the author believes that a detailed study of such projections is needed, and this was begun by Boca [2] in his construction of Fourier invariant projections, and later by the author in [18] where projections orthogonal to their Fourier transform are constructed (such projections are needed in constructing direct sums of two identical circle algebras in which the Fourier transform swaps the two summands). In [17], a clue is given of the inductive

1991 *Mathematics Subject Classification.* 46L80, 46L40, 46L35.

Key words and phrases. C*-algebras, irrational rotation algebras, automorphisms, inductive limits, K-groups, AF-algebras, Theta functions.

Research partly supported by NSERC grant OGP0169928

(TeXFile: pointproj.tex)

limit structure of the Fourier transform by the construction of a (noncanonical) model of an inductive limit automorphism of order four on A_θ that agrees with σ on K_1 (a pseudo-Fourier transform) such that the fixed point subalgebra is an AF-algebra. In [15] and [16], it was shown for the crossed product $A_\theta \rtimes_\sigma \mathbb{Z}_4$ that there is always an inclusion $\mathbb{Z}^9 \hookrightarrow K_0$, which, for a dense G_δ set of θ 's, is an isomorphism (and $K_1 = 0$). The main thrust of this paper is to begin tackling Elliott's problem more directly (and canonically) by showing that one can construct infinitely many partially approximating finite dimensional subalgebras of A_θ (i.e., points) that are invariant under the Fourier transform, as well as invariant direct sums of two circle subalgebras. More precisely, the following result is proved.

Main Theorem. *Let θ be a transcendental number. Then the irrational rotation algebra A_θ contains infinitely many C^* -subalgebras $\mathcal{M}_q^2 = \mathcal{M}_q \oplus \mathcal{M}_q$ (orthogonal sum), where $\mathcal{M}_q$ is isomorphic to $M_q(\mathbb{C})$ and q is a perfect square, such that*

- (1) *the Fourier transform maps each summand of $\mathcal{M}_q^2$ onto the other,*
- (2) *the unit projection e of $\mathcal{M}_q$ is approximately central,*
- (3) *the compressions eUe and eVe are approximately contained in $\mathcal{M}_q$,*
- (4) *the projection e is also the unit of a subalgebra C isomorphic to $M_q(C(\mathbb{T}))$ such that $C \oplus \sigma(C)$ is a Fourier invariant circle algebra,*

Section 2 contains the background and notation used in later sections. In Section 3 the construction of $\mathcal{M}_q$ is carried out to give (1) and (4). In Sections 4 and 5 we prove (3) and (2), respectively. Section 6 is an Appendix containing some basic lemmas used in the approximations of Sections 4 and 5.

Throughout, we shall adopt the notation $e(t) := e^{2\pi it}$.

§2. Initial Considerations

In order to focus on the C^* -algebraic aspects of the problem, and to avoid some number theoretic obstacles, we shall from now on assume that θ is a transcendental number in $(0, 1)$. This will simplify the choice of the lattice D in Section 3. The following result is an application of the Thue-Siegel-Roth Theorem (see for example [9], Chapter 7).

Proposition 2.1. *Let θ be a transcendental number in $(0, 1)$. For each $a > 0$, there are infinitely many convergents p/q such that $p, q, q - p$ are all perfect squares and satisfying*

$$\left| \theta - \frac{p}{q} \right| < \frac{1}{q^a}. \quad (2.1)$$

Proof. Since θ is transcendental, so is

$$\xi = \frac{1 - \sqrt{\theta}}{\sqrt{1 - \theta}}$$

which is in $(0, 1)$. Thanks to the Thue-Siegel-Roth Theorem, for each positive number ϵ there are infinitely many convergents $r/s \in (0, 1)$ of θ that satisfy

$$\left| \frac{r}{s} - \xi \right| < \frac{1}{s^\epsilon}.$$

Let $f(x) = 2x/(x^2 + 1)$ so that $f(\xi) = \sqrt{1 - \theta}$. One clearly has

$$|f(x) - f(y)| = \frac{2|x - y||1 - xy|}{(x^2 + 1)(y^2 + 1)}. \quad (2.2)$$

Let $m = 2rs$, $k = s^2 - r^2$, $n = r^2 + s^2$, so that $m^2 + k^2 = n^2$ and $\frac{m}{n} = f(\frac{r}{s})$. Since ξ and r/s are in $(0, 1)$, from (2.2) we have

$$\left| \frac{m}{n} - \sqrt{1 - \theta} \right| = |f(\frac{r}{s}) - f(\xi)| < 2 \left| \frac{r}{s} - \xi \right| < \frac{2}{s^\epsilon}.$$

One gets

$$\left| \theta - \frac{n^2 - m^2}{n^2} \right| = \left| \frac{m^2}{n^2} - (1 - \theta) \right| < \frac{4}{s^\epsilon}.$$

Now since $r < s$, we have

$$\frac{4}{s^\epsilon} = \frac{4 \cdot 2^{\epsilon/2}}{(2s^2)^{\epsilon/2}} < \frac{4 \cdot 2^{\epsilon/2}}{(r^2 + s^2)^{\epsilon/2}} = \frac{4 \cdot 2^{\epsilon/2}}{n^{\epsilon/2}}.$$

Therefore, taking $p = n^2 - m^2 = k^2$ and $q = n^2$ (so $q - p = m^2$) they satisfy

$$\left| \theta - \frac{p}{q} \right| < \frac{4 \cdot 2^{\epsilon/2}}{n^{\epsilon/2}} = \frac{4 \cdot 2^{\epsilon/2}}{q^{\epsilon/4}} < \frac{1}{q^a}$$

where the last inequality holds, for example by taking $\epsilon = 4(1 + a)$ and for large enough q . Since the greatest common divisor d of p and q is also a perfect square, $\frac{p}{d}, \frac{q}{d}, \frac{q-p}{d}$ are perfect squares and satisfy the preceding inequality, so one can assume p, q are relatively prime. In this case, it is well-known (for $q > 2$) that (2.1) in itself implies that p/q must be a convergent of θ (of course, providing that $a \geq 3$). $\square$

If θ is algebraic, then by Roth's Theorem [13], for any $\epsilon > 0$ the inequality $|\theta - \frac{p}{q}| < \frac{1}{q^{2+\epsilon}}$ can only be satisfied by *finitely* many rationals p/q , a result for which he received the 1958 Fields Medal. Therefore, our approach here only applies to transcendental numbers θ , and throughout the paper we shall let p/q be a fixed convergent of θ such that p, q , and $q - p$ are perfect squares satisfying (2.1) with $a = 3$:

$$\left| \theta - \frac{p}{q} \right| < \frac{1}{q^3}.$$

Further, without loss of generality, we may assume that there are infinitely many such convergents such that $\frac{p}{q} < \theta$ (otherwise, one replaces θ by $1 - \theta$). (If there are infinitely many convergents p/q such that $\theta < \frac{p}{q}$ and $p, q, q - p$ are perfect squares, then $\frac{q-p}{q}$ is also a convergent with $\frac{q-p}{q} < 1 - \theta$ where $q - p$ is a perfect square.)

Theta Functions. The main Theta functions used in this paper are

$$\vartheta_2(z, t) = \sum_n e^{\pi i t (n + \frac{1}{2})^2} e^{i 2 z (n + \frac{1}{2})}, \quad \vartheta_3(z, t) = \sum_n e^{\pi i t n^2} e^{i 2 z n}$$

for $z, t \in \mathbb{C}$ and $\text{Im}(t) > 0$, where all summations range over the integers. (For a classic treatment see [19].)

A Gaussian Theta Function. In [18] a Schwartz function h on $\mathbb{R}$ was constructed that lead to the existence of flip invariant projections e in A_θ orthogonal to their Fourier transform (i.e., $e\sigma(e) = 0$). Since this function is important for the present paper, we shall recall briefly some of its properties. Let $\alpha > 0, \beta > 2$ be such that $\beta^2 = 4(\alpha^2 + 1)$. Define

$$h(x) = e^{-\pi\alpha x^2} \sum_p e^{-\pi\alpha p^2} e^{\pi\alpha p} e([\frac{\beta}{2}p - \frac{\beta}{4}]x)$$

an even real-valued Schwartz function on $\mathbb{R}$. It was shown in [18] that

$$\int_{\mathbb{R}} \overline{h(x)} \widehat{h}(x + \beta m) e(\beta n x) dx = 0 \quad (2.3)$$

for all integers m, n , where $\widehat{h}$ is the usual Fourier transform of h (defined below). It was also shown that for all real s, t one has

$$H(s, t) := \int_{\mathbb{R}} \overline{h(x)} h(x + s) e(xt) dx = \frac{1}{\sqrt{2\alpha}} e(-\frac{st}{2}) e^{-\pi\alpha s^2/2} e^{-\pi t^2/(2\alpha)} \Gamma(\frac{t}{\beta}, \frac{s}{\beta}) \quad (2.4)$$

where

$$\Gamma(u, v) = e^{\pi\alpha/2} \left[\vartheta_2(\frac{\pi}{2}\beta^2 v, 2i\alpha) \vartheta_3(i\frac{\pi}{2\alpha}\beta^2 u, it_\alpha) + \vartheta_3(\frac{\pi}{2}\beta^2 v, 2i\alpha) \vartheta_2(i\frac{\pi}{2\alpha}\beta^2 u, it_\alpha) \right] \quad (2.5)$$

for $u, v \in \mathbb{R}$, and $t_\alpha = 4\alpha + \frac{2}{\alpha}$ ([18, Lemma 4.2]). It is clear that Γ is even in each variable and is real valued (in fact all of its terms are positive, except possibly $\vartheta_2(\frac{\pi}{2}\beta^2 v, 2i\alpha)$). Using the fact that

$$2 < e^{\pi\alpha/2} \vartheta_2(0, 2i\alpha) < 2\vartheta_3(0, 2i\alpha) \quad (2.6)$$

one has

$$|\Gamma(u, v)| \leq \vartheta_3(0, 2i\alpha) \left[2\vartheta_3(i\frac{\pi}{2\alpha}\beta^2 u, it_\alpha) + e^{\pi\alpha/2} \vartheta_2(i\frac{\pi}{2\alpha}\beta^2 u, it_\alpha) \right]. \quad (2.7)$$

We also had the period 1 functions

$$\psi_n(t) := \sum_m e(\beta^2 mn/2) e^{-\pi\beta^2 m^2/(2\alpha)} \Gamma(m, n) e(mt) \quad (2.8)$$

for $n \in \mathbb{Z}$, which will arise in the cut down approximations below. The function ψ_0 is positive invertible and it was shown that we have the inequality

$$\frac{|\psi_n(t)|}{\psi_0(s)} \leq 1.21 \frac{\vartheta_3(0, 2i\alpha)}{\vartheta_3(\frac{\pi}{2}, 2i\alpha)}.$$

for all real s, t and any integer n ([18, Lemma 4.3]). The right side of this inequality is a decreasing function of α and hence is no more than its value at $\alpha = \frac{1}{4}$, hence

$$\frac{|\psi_n(t)|}{\psi_0(s)} \leq 1.21 \frac{\vartheta_3(0, \frac{i}{2})}{\vartheta_3(\frac{\pi}{2}, \frac{i}{2})} = 1.21(1 + \sqrt{2}) < 3 \quad (2.9)$$

for $\alpha > \frac{1}{4}$. Further, for $\beta > \beta_0 := 2.063$ and all real t one has

$$1 < \psi_0(t) < 5, \quad \text{and} \quad \psi_0(t) \rightarrow 2 \text{ uniformly in } t \text{ as } \beta \rightarrow \infty. \quad (2.10)$$

([18, Proposition 4.4]).

Rieffel's Framework. We briefly sketch Rieffel's setup in [12] and the notation to be used. Let M be a locally compact Abelian group and let $G = M \times \hat{M}$, where $\hat{M}$ is the dual of M . Let $\mathfrak{h}$ denote the Heisenberg cocycle on the group $G = M \times \hat{M}$ given by $\mathfrak{h}((m, s), (m', s')) = \langle m, s' \rangle$, where $\langle m, s' \rangle$ is the canonical pairing $M \times \hat{M} \rightarrow \mathbb{T}$. The Heisenberg (unitary) representation $\pi : G \rightarrow \mathcal{U}(L^2(M))$ of G is given by

$$(\pi_{(m,s)}f)(n) = \langle n, s \rangle f(n + m)$$

where $m, n \in M$, $s \in \hat{M}$, and $f \in L^2(M)$. It has the properties

$$\pi_x \pi_y = \mathfrak{h}(x, y) \pi_{x+y} = \mathfrak{h}(x, y) \overline{\mathfrak{h}(y, x)} \pi_y \pi_x, \quad \pi_x^* = \mathfrak{h}(x, x) \pi_{-x}$$

for $x, y \in G$. If D is a given lattice in G (i.e., discrete subgroup such that G/D is compact), its covolume $|G/D|$ is the Haar measure of a fundamental domain for D in G . Its associated twisted group C*-algebra $C^*(D, \mathfrak{h})$ is the C*-subalgebra of the bounded operators on $L^2(M)$ generated by the unitaries π_x for $x \in D$. The complementary lattice of D is

$$D^\perp = \{y \in G : \mathfrak{h}(x, y) \overline{\mathfrak{h}(y, x)} = 1, \forall x \in D\}.$$

The C*-algebra $C^*(D^\perp, \bar{\mathfrak{h}})$ can be viewed as the C*-subalgebra of bounded operators on $L^2(M)$ generated by the unitaries π_y^* for $y \in D^\perp$. By Rieffel's Theorem 2.15 [12], the Schwartz space on M , denoted $\mathcal{S}(M)$, is an equivalence bimodule with $C^*(D, \mathfrak{h})$ acting on the left and $C^*(D^\perp, \bar{\mathfrak{h}})$ acting on the right by

$$\begin{aligned} af &= \int_D a(x) \pi_x(f) dx = |G/D| \sum_{x \in D} a(x) \pi_x(f) \\ fb &= \int_{D^\perp} b(y) \pi_y^*(f) dy = \sum_{y \in D^\perp} b(y) \pi_y^*(f) \end{aligned}$$

where $f \in \mathcal{S}(M)$, $a \in C^*(D, \mathfrak{h})$, $b \in C^*(D^\perp, \bar{\mathfrak{h}})$, and where the mass point measure (dx) on D is $|G/D|$ and on $D^\perp$ it is one. The inner products on $\mathcal{S}(M)$ with values in the algebras $C^*(D, \mathfrak{h})$ and $C^*(D^\perp, \bar{\mathfrak{h}})$ are given, respectively, by

$$\langle f, g \rangle_D = |G/D| \sum_{w \in D} \langle f, g \rangle_D(w) \pi_w, \quad \langle f, g \rangle_{D^\perp} = \sum_{z \in D^\perp} \langle f, g \rangle_{D^\perp}(z) \pi_z^*$$

where

$$\begin{aligned} \langle f, g \rangle_D(w_1, w_2) &= \int_M f(x) \overline{g(x + w_1)} \langle x, w_2 \rangle dx \\ \langle f, g \rangle_{D^\perp}(z_1, z_2) &= \int_M \overline{f(x)} g(x + z_1) \langle x, z_2 \rangle dx \end{aligned}$$

where $(w_1, w_2) \in D$ and $(z_1, z_2) \in D^\perp$. These satisfy the associativity condition

$$\langle f, g \rangle_D h = f \langle g, h \rangle_{D^\perp}.$$

(See [12], pages 266 and 269.) The canonical normalized traces are given by

$$\tau_D \left(\sum_{w \in D} a_w \pi_w \right) = a_0, \quad \tau_{D^\perp} \left(\sum_{z \in D^\perp} b_z \pi_z^* \right) = b_0,$$

($a_w, b_w \in \mathbb{C}$) and satisfy the equality

$$\tau_D(\langle f, g \rangle_D) = |G/D| \tau_{D^\perp}(\langle g, f \rangle_{D^\perp}).$$

From this it follows that if ξ is a Schwartz function such that $\langle \xi, \xi \rangle_{D^\perp} = 1$, then $e = \langle \xi, \xi \rangle_D$ is a projection in $C^*(D, \mathfrak{h})$ of trace $|G/D|$. In this case, one has the cut down isomorphism

$$\mu_\xi : eC^*(D, \mathfrak{h})e \rightarrow C^*(D^\perp, \bar{\mathfrak{h}})$$

given by

$$\mu_\xi(xex) = \langle \xi, x\xi \rangle_{D^\perp}, \quad \mu_\xi^{-1}(y) = \langle \xi y, \xi \rangle_D$$

for $x \in C^*(D, \mathfrak{h})$ and $y \in C^*(D^\perp, \bar{\mathfrak{h}})$. Projections e arising in this manner might be called *generalized Rieffel projections*, since in [7], Elliott and Lin showed that the classical Rieffel projections [11] in the rotation algebra do in fact have this inner product form.

In the present paper, we shall take $M = \mathbb{R} \times \mathbb{F}$ where $\mathbb{F}$ is a finite abelian group so that $\hat{M} = M$ in a natural fashion. With this identification in mind one is permitted to define the order four automorphism $R : G \rightarrow G$ by $R(u; v) = (-v; u)$, $u, v \in M$. If D is a lattice subgroup of G such that $R(D) = D$ (and hence $R(D^\perp) = D^\perp$), then (as in [2]) there are order four automorphisms σ, σ' of $C^*(D, \mathfrak{h})$ and $C^*(D^\perp, \bar{\mathfrak{h}})$, respectively, that satisfy

$$\sigma(\pi_w) = \overline{\mathfrak{h}(w, w)} \pi_{Rw}, \quad \sigma'(\pi_z^*) = \mathfrak{h}(z, z) \pi_{Rz}^*,$$

for $w \in D$, $z \in D^\perp$, and

$$\sigma(\langle f, g \rangle_D) = \langle \hat{f}, \hat{g} \rangle_D, \quad \sigma'(\langle f, g \rangle_{D^\perp}) = \langle \hat{f}, \hat{g} \rangle_{D^\perp},$$

where $\hat{f}$ is the usual Fourier transform of $f \in \mathcal{S}(M)$ given by

$$\hat{f}(s) = \int_M f(x) \overline{\langle x, s \rangle} dx$$

for $s \in M$. Note that the isomorphism μ_ξ commutes with the Fourier transform on each algebra according to the commutative diagram

$$\begin{array}{ccc} eC^*(D)e & \xrightarrow{\mu_\xi} & C^*(D^\perp) \\ \sigma \downarrow & & \downarrow \sigma' \\ \sigma(e)C^*(D)\sigma(e) & \xrightarrow{\mu_{\hat{\xi}}} & C^*(D^\perp) \end{array}$$

which, for $x \in eC^*(D)e$, follows from

$$\sigma'(\mu_\xi(x)) = \sigma'(\langle \xi, x\xi \rangle_{D^\perp}) = \langle \hat{\xi}, \widehat{x\xi} \rangle_{D^\perp} = \langle \hat{\xi}, \sigma(x)\hat{\xi} \rangle_{D^\perp} = \mu_{\hat{\xi}}(\sigma(x))$$

as $\sigma(e) = \langle \hat{\xi}, \hat{\xi} \rangle_D$. As in [7], we shall write “ xex ” := $\langle \xi, x\xi \rangle_{D^\perp}$.

§3. The Construction

Let $M = \mathbb{R} \times \mathbb{Z}_q \times \mathbb{Z}_q$, and consider the lattice D in $G = M \times \hat{M}$ with basis

$$D : \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \end{bmatrix} = \begin{bmatrix} a & [p']_q & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a & [p']_q & 0 \end{bmatrix},$$

where p' is a positive integer such that $p = (p')^2$, where we recall that $\frac{p}{q} < \theta$, and where $a = (\theta - \frac{p}{q})^{1/2}$. This lattice is clearly invariant under $R : (w_1, w_2) \mapsto (-w_2, w_1)$. The algebra $C^*(D, \mathfrak{h})$ is generated by the canonical unitaries $U_1 = \pi_{\varepsilon_1}$, $U_2 = \pi_{\varepsilon_2}$ whose commutation relation is

$$U_1 U_2 U_1^* U_2^* = \pi_{\varepsilon_1} \pi_{\varepsilon_2} \pi_{\varepsilon_1}^* \pi_{\varepsilon_2}^* = \mathfrak{h} \mathfrak{h}^*(\varepsilon_1, \varepsilon_2) = e(a^2 + \frac{(p')^2}{q}) = e(\theta)$$

Therefore, U_1, U_2 generate the irrational rotation algebra A_θ , and the Fourier transform enjoys $\sigma(U_1) = U_2$, $\sigma(U_2) = U_1^{-1}$. A fundamental domain for D is $[0, a) \times \mathbb{Z}_q \times \mathbb{Z}_q \times [0, a) \times \mathbb{Z}_q \times \mathbb{Z}_q$ so its covolume is $|G/D| = a^2 q^2 = q(q\theta - p) < 1$. Put

$$\beta := \frac{1}{qa}.$$

Observe that since by hypothesis $a^2 < q^{-3}$, one has $\beta > \sqrt{q}$ so that $\beta > 2$ for $q \geq 4$. The annihilator lattice $D^\perp$ can be checked to be spanned by the basis elements

$$D^\perp : \begin{bmatrix} \delta_1 \\ \delta_2 \\ \delta_3 \\ \delta_4 \end{bmatrix} = \begin{bmatrix} \beta & [-p_0] & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \beta & [-p_0] & 0 \\ 0 & 0 & [1] & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & [1] \end{bmatrix}$$

where p_0 is an integer such that $p_0 p' = 1 \pmod{q}$. Let $V_j = \pi_{\delta_j}^* = \pi_{-\delta_j}$. Then

$$V_1 V_2 = e(\mu) V_2 V_1, \quad V_3 V_4 = e(\frac{1}{q}) V_4 V_3, \quad V_j V_k = V_k V_j$$

for $j = 1, 2$ and $k = 3, 4$, where $\mu = \beta^2 + \frac{p^2}{q}$ is in the $\text{GL}(2, \mathbb{Z})$ orbit of θ . One has $C^*(D^\perp, \bar{\mathfrak{h}}) = C^*(V_j) \cong M_q(A_\mu)$.

The $D^\perp$ inner product becomes, for $f, g \in \mathcal{S}(M)$,

$$\begin{aligned} \langle f, g \rangle_{D^\perp} &= \langle f, g \rangle_{D^\perp} (n_1 \delta_1 + n_2 \delta_2 + n_3 \delta_3 + n_4 \delta_4) \\ &= \langle f, g \rangle_{D^\perp} (n_1 \beta, [-n_1 p_0], [n_3]; n_2 \beta, [-n_2 p_0], [n_4]) \\ &= \int_{\mathbb{R} \times \mathbb{Z}_q \times \mathbb{Z}_q} \overline{f(x, [n], [m])} g(x + n_1 \beta, [n - n_1 p_0], [m + n_3]) e(x n_2 \beta - \frac{n n_2 p_0}{q} + \frac{m n_4}{q}) dx d[n] d[m] \\ &= \frac{1}{q} \sum_{[n], [m] \in \mathbb{Z}_q} e(-\frac{n n_2 p_0}{q} + \frac{m n_4}{q}) \int_{\mathbb{R}} \overline{f(x, [n], [m])} g(x + n_1 \beta, [n - n_1 p_0], [m + n_3]) e(x n_2 \beta) dx. \end{aligned}$$

Since

$$\pi_{n_1 \delta_1 + n_2 \delta_2 + n_3 \delta_3 + n_4 \delta_4}^* = V_1^{n_1} V_2^{n_2} V_3^{n_3} V_4^{n_4}$$

one has (where $n_3, n_4 = 0, 1, \dots, q-1$ and $n_1, n_2 \in \mathbb{Z}$)

$$\langle f, g \rangle_{D^\perp} = \sum_{n_1 n_2 n_3 n_4} \langle f, g \rangle_{D^\perp}(\Sigma n_* \delta_*) \cdot V_1^{n_1} V_2^{n_2} V_3^{n_3} V_4^{n_4}.$$

Now suppose that f and g are Schwartz functions on M of the form

$$f(x, [n], [m]) = h(x) \varphi_f([n], [m]), \quad g(x, [n], [m]) = h'(x) \varphi_g([n], [m])$$

where $h, h' \in \mathcal{S}(\mathbb{R})$ and φ_f, φ_g are any functions. Then one gets

$$\langle f, g \rangle_{D^\perp}(n_1 \delta_1 + n_2 \delta_2 + n_3 \delta_3 + n_4 \delta_4) = \frac{1}{q} \Omega_{n_1 n_2 n_3 n_4}(\varphi_f, \varphi_g) \cdot [hh'](n_1 \beta, n_2 \beta)$$

where

$$\Omega_{n_1 n_2 n_3 n_4}(\varphi_f, \varphi_g) = \sum_{[n], [m] \in \mathbb{Z}_q} e(-\frac{nn_2 p_0}{q} + \frac{mn_4}{q}) \overline{\varphi_f([n], [m])} \varphi_g([n - n_1 p_0], [m + n_3])$$

and

$$[hh'](s, t) = \int_{\mathbb{R}} \overline{h(x)} h'(x + s) e(xt) dx.$$

First, observe that if h is chosen so that $[h\hat{h}](n_1, n_2) = 0$ for all integers n_1, n_2 , then $\langle f, \hat{f} \rangle_{D^\perp} = 0$. (This follows from $\hat{f}(x, [n], [m]) = \hat{h}(x) \widehat{\varphi_f}([n], [m])$.) Consider the Schwartz function

$$f(x, [n], [m]) = (2\alpha)^{1/4} h(x) \delta_q^{n-m}$$

where $\delta_q^k = 1$ if $q|k$ and 0 otherwise, h is the Gaussian Theta function of Section 2, and $\alpha > 0$ satisfies $\beta^2 = 4(\alpha^2 + 1)$. We have

$$\begin{aligned} \Omega_{n_1 n_2 n_3 n_4}(\varphi_f, \varphi_f) &= \sum_{m, n=0}^{q-1} e(-\frac{nn_2 p_0}{q} + \frac{mn_4}{q}) \delta_q^{n-m} \delta_q^{n-m-n_1 p_0 - n_3} \\ &= \delta_q^{n_1 p_0 + n_3} \sum_{n=0}^{q-1} e(-\frac{n}{q}(n_2 p_0 - n_4)) \\ &= q \delta_q^{n_1 p_0 + n_3} \delta_q^{n_2 p_0 - n_4} \end{aligned}$$

hence (with $n_3, n_4 = 0, 1, \dots, q-1$ and $n_1, n_2 \in \mathbb{Z}$)

$$\begin{aligned} \langle f, f \rangle_{D^\perp} &= \frac{\sqrt{2\alpha}}{q} \sum_{n_1 n_2 n_3 n_4} q \delta_q^{n_1 p_0 + n_3} \delta_q^{n_2 p_0 - n_4} H(n_1 \beta, n_2 \beta) V_1^{n_1} V_2^{n_2} V_3^{n_3} V_4^{n_4} \\ &= \sqrt{2\alpha} \sum_{n_1 n_2} H(n_1 \beta, n_2 \beta) V_1^{n_1} V_2^{n_2} V_3^{-n_1 p_0} V_4^{n_2 p_0} \\ &=: b \end{aligned}$$

where (since the pair V_1, V_2 commutes with V_3, V_4)

$$b = \sqrt{2\alpha} \sum_{n_1 n_2} H(n_1 \beta, n_2 \beta) W_1^{n_1} W_2^{n_2}, \quad W_1 = V_1 V_3^{-p_0}, \quad W_2 = V_2 V_4^{p_0}$$

and $W_1 W_2 = e(\beta^2) W_2 W_1$. (Recall that H is given by (2.4).) The following lemma shows that b is an invertible element (at least) for sufficiently large q .

Lemma 3.1. *One has $\|b - \psi_0\| \rightarrow 0$ as $q \rightarrow \infty$ and hence $\|b - 2\| \rightarrow 0$. In particular, b is invertible for sufficiently large q .*

Proof. Since

$$H(m\beta, n\beta) = \frac{1}{\sqrt{2\alpha}} e(-\frac{1}{2}\beta^2 mn) e^{-\pi\alpha\beta^2 m^2/2} e^{-\pi\beta^2 n^2/(2\alpha)} \Gamma(n, m)$$

we can write

$$b = \sum_m e^{-\pi\alpha\beta^2 m^2/2} W_1^m \rho_m$$

where $\rho_m = \rho_m(W_2)$ can be viewed as a period one function as

$$\rho_m(t) = \sum_n e(-\frac{1}{2}\beta^2 mn) e^{-\pi\beta^2 n^2/(2\alpha)} \Gamma(n, m) e(nt) = \psi_m(t - \beta^2 m)$$

where ψ_m is given by (2.8). In particular, $\rho_0 = \psi_0$ is strictly positive. From (2.9) one has $\|\rho_m \rho_0^{-1}\| < 3$ for all m (and $\alpha > \frac{1}{4}$), hence one obtains

$$\|b\psi_0^{-1} - I\| \leq \sum_{m \neq 0} e^{-\pi\alpha\beta^2 m^2/2} \|\rho_m \rho_0^{-1}\| < 3 \sum_{m \neq 0} e^{-\pi\alpha\beta^2 m^2/2} = 3[\vartheta_3(0, \frac{i}{2}\alpha\beta^2) - 1] =: C_q$$

and $C_q \rightarrow 0$ as $q \rightarrow \infty$. Thus one gets $\|b - \psi_0\| < C_q \|\psi_0\| < 5C_q$. Since $\|\psi_0 - 2\| \rightarrow 0$ as $\beta \rightarrow \infty$ by (2.10), one deduces that $\|b - 2\| \rightarrow 0$. $\square$

Now since $\|x - 1\| < \epsilon < 1$ implies $\|x^{-1} - 1\| < \epsilon/(1 - \epsilon)$, one gets (for large enough q)

$$\|\psi_0 b^{-1} - I\| < \frac{C_q}{1 - C_q}$$

and using (2.10)

$$\|b^{-1} - \psi_0^{-1}\| < \frac{C_q}{1 - C_q} \|\psi_0^{-1}\| < \frac{C_q}{1 - C_q}.$$

This proves the following lemma to be used below.

Lemma 3.2. *The norms $\|b\|$, $\|b^{-1/2}\|$ are uniformly bounded for $\alpha > \frac{1}{4}$.*

Putting $\xi = fb^{-1/2}$ one has $\langle \xi, \xi \rangle_{D^\perp} = 1$ and hence $e = \langle \xi, \xi \rangle_D$ is a projection in $C^*(D, \mathfrak{h}) = A_\theta$ with trace $\tau(e) = |G/D| = q(q\theta - p)$. Since b is flip invariant, because the flip $(V_j \mapsto V_j^{-1})$ maps W_j to W_j^{-1} and $H(-m\beta, -n\beta) = H(m\beta, n\beta)$, it follows that e is a flip invariant projection in A_θ such that $e\sigma(e) = 0$ (by (2.3)). We conclude that e is the unit projection of a matrix subalgebra $\mathcal{M}_q$ of A_θ obtained from the Morita isomorphism μ_ξ at the end of Section 2. Hence A_θ contains the finite dimensional C^* -subalgebra $\mathcal{M}_q^2 = \mathcal{M}_q \oplus \sigma(\mathcal{M}_q)$ in which its summands are mapped onto one another by the Fourier transform (and whose unit $e + \sigma(e)$ is Fourier invariant). If one looks at the C^* -algebra generated by V_1, V_3, V_4 , which is a $q \times q$ matrix algebra over $C(\mathbb{T})$, and transforms it by μ_ξ , one obtains a circle subalgebra of A_θ whose unit is also e . This proves (1) and (4) of the Main Theorem. Since from the following two sections e is shown to be approximately central, and that the compressions eU_1e, eU_2e are approximately contained in $\mathcal{M}_q$, it follows that $\mathcal{M}_q^2$ is a partially approximating C^* -subalgebra of A_θ that is invariant under the Fourier transform.

Remark. It is immediate from the present construction that one also obtains Fourier orthogonal projections of trace $k(q\theta - p)$ for $k = 1, \dots, q$. (By looking at subprojections of e .)

§4. Cut Down Approximation

In this section the compressions eU_1e and eU_2e are approximated by elements of the matrix algebra generated by V_3, V_4 of the previous section. More precisely:

Proposition 4.1. *One has*

$$\|\langle \xi, U_1 \xi \rangle_{D^\perp} - V_3^{p'}\| \rightarrow 0, \quad \|\langle \xi, U_2 \xi \rangle_{D^\perp} - V_4^{-p'}\| \rightarrow 0,$$

as $q \rightarrow \infty$.

Since $U_1 = \pi_{\varepsilon_1}$,

$$(U_1 f)(x, [r], [s]) = f(x + a, [r + p'], [s])$$

and thus

$$\begin{aligned} & \langle f, U_1 f \rangle_{D^\perp} (\Sigma n_* \delta_*) \\ &= \frac{1}{q} \sum_{m,n=0}^{q-1} e(-\frac{nn_2 p_0}{q} + \frac{mn_4}{q}) \int_{\mathbb{R}} \overline{f(x, [n], [m])} (U_1 f)(x + n_1 \beta, [n - n_1 p_0], [m + n_3]) e(x n_2 \beta) dx \\ &= \frac{(2\alpha)^{\frac{1}{4}}}{q} \sum_{m,n=0}^{q-1} e(-\frac{nn_2 p_0}{q} + \frac{mn_4}{q}) \int_{\mathbb{R}} \overline{h(x)} \delta_q^{n-m} f(x + n_1 \beta + a, [n - n_1 p_0 + p'], [m + n_3]) e(x n_2 \beta) dx \\ &= \frac{\sqrt{2\alpha}}{q} \sum_{m,n=0}^{q-1} e(-\frac{nn_2 p_0}{q} + \frac{mn_4}{q}) \delta_q^{n-m} \delta_q^{n-m-n_1 p_0 + p' - n_3} \int_{\mathbb{R}} \overline{h(x)} h(x + n_1 \beta + a) e(x n_2 \beta) dx \\ &= \frac{\sqrt{2\alpha}}{q} \delta_q^{n_1 p_0 - p' + n_3} \sum_{n=0}^{q-1} e(-\frac{n}{q} (n_2 p_0 - n_4)) H(n_1 \beta + a, n_2 \beta) \\ &= \sqrt{2\alpha} \delta_q^{n_1 p_0 - p' + n_3} \delta_q^{n_2 p_0 - n_4} H(n_1 \beta + a, n_2 \beta). \end{aligned}$$

Therefore,

$$\begin{aligned} \langle f, U_1 f \rangle_{D^\perp} &= \sqrt{2\alpha} \sum_{n_1 n_2 n_3 n_4} \delta_q^{n_1 p_0 - p' + n_3} \delta_q^{n_2 p_0 - n_4} H(n_1 \beta + a, n_2 \beta) V_1^{n_1} V_2^{n_2} V_3^{n_3} V_4^{n_4} \\ &= \sqrt{2\alpha} \sum_{n_1 n_2} H(n_1 \beta + a, n_2 \beta) V_1^{n_1} V_2^{n_2} V_3^{p' - n_1 p_0} V_4^{n_2 p_0} \\ &= V_3^{p'} X \end{aligned}$$

where

$$X = \sqrt{2\alpha} \sum_{m,n} H(m\beta + a, n\beta) W_1^m W_2^n$$

is a perturbation of the element b of Section 3. This gives

$$“eU_1e” = \langle \xi, U_1 \xi \rangle_{D^\perp} = b^{-1/2} V_3^{p'} X b^{-1/2}.$$

One similarly calculates $\langle f, U_2 f \rangle_{D^\perp}$. Since $U_2 = \pi_{\varepsilon_2}$,

$$(U_2 f)(x, r, s) = e(ax + \frac{rp'}{q}) f(x, r, s)$$

we have

$$\begin{aligned}
& \langle f, U_2 f \rangle_{D^\perp} (\Sigma n_* \delta_*) \\
&= \frac{1}{q} \sum_{m,n=0}^{q-1} e(-\frac{nn_2 p_0}{q} + \frac{mn_4}{q}) \int_{\mathbb{R}} \overline{f(x, [n], [m])} (U_2 f)(x + n_1 \beta, [n - n_1 p_0], [m + n_3]) e(x n_2 \beta) dx \\
&= \frac{\sqrt{2\alpha}}{q} e(an_1 \beta) \sum_{m,n=0}^{q-1} e(-\frac{nn_2 p_0}{q} + \frac{mn_4}{q}) e(\frac{(n-n_1 p_0)p'}{q}) \delta_q^{n-m} \delta_q^{n-m-n_1 p_0-n_3} H(n_1 \beta, n_2 \beta + a) \\
&= \frac{\sqrt{2\alpha}}{q} e(\frac{n_1}{q}) e(\frac{-n_1 p_0 p'}{q}) \sum_{m,n=0}^{q-1} e(-\frac{nn_2 p_0}{q} + \frac{mn_4}{q}) e(\frac{np'}{q}) \delta_q^{n-m} \delta_q^{n-m-n_1 p_0-n_3} H(n_1 \beta, n_2 \beta + a) \\
&= \frac{\sqrt{2\alpha}}{q} \delta_q^{n_1 p_0 + n_3} \sum_{n=0}^{q-1} e(-\frac{nn_2 p_0}{q} + \frac{nn_4}{q} + \frac{np'}{q}) H(n_1 \beta, n_2 \beta + a) \\
&= \frac{\sqrt{2\alpha}}{q} \delta_q^{n_1 p_0 + n_3} \sum_{n=0}^{q-1} e(-\frac{n}{q} (n_2 p_0 - n_4 - p')) H(n_1 \beta, n_2 \beta + a) \\
&= \sqrt{2\alpha} \delta_q^{n_1 p_0 + n_3} \delta_q^{n_2 p_0 - n_4 - p'} H(n_1 \beta, n_2 \beta + a)
\end{aligned}$$

hence

$$\begin{aligned}
\langle f, U_2 f \rangle_{D^\perp} &= \sqrt{2\alpha} \sum_{n_1 n_2 n_3 n_4} \delta_q^{n_1 p_0 + n_3} \delta_q^{n_2 p_0 - n_4 - p'} H(n_1 \beta, n_2 \beta + a) V_1^{n_1} V_2^{n_2} V_3^{n_3} V_4^{n_4} \\
&= \sqrt{2\alpha} \sum_{n_1 n_2} H(n_1 \beta, n_2 \beta + a) V_1^{n_1} V_2^{n_2} V_3^{-n_1 p_0} V_4^{n_2 p_0 - p'} \\
&= Y V_4^{-p'}
\end{aligned}$$

where

$$Y = \sqrt{2\alpha} \sum_{mn} H(m\beta, n\beta + a) W_1^m W_2^n.$$

Thus

$$"eU_2 e" = \langle \xi, U_2 \xi \rangle_{D^\perp} = b^{-1/2} Y V_4^{-p'} b^{-1/2}.$$

Lemma 4.2. *The norms $\|X - b\|$ and $\|Y - b\|$ converge to zero as $q \rightarrow \infty$.*

Proof. We shall prove this for X only, the proof for Y being similar. We have

$$\begin{aligned}
\|X - b\| &\leq \sqrt{2\alpha} \sum_{m,n} |H(m\beta + a, n\beta) - H(m\beta, n\beta)| \\
&= \sum_{m,n} e^{-\frac{\pi\beta^2 n^2}{2\alpha}} e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} \left| e(-\frac{1}{2}n\beta a) e^{-\frac{1}{2}\pi\alpha a^2} e^{-\pi\alpha\beta am} \Gamma(n, m + \frac{a}{\beta}) - \Gamma(n, m) \right| \\
&= A + 2B
\end{aligned}$$

where

$$A = \sum_m e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} \left| e^{-\frac{1}{2}\pi\alpha a^2} e^{-\pi\alpha\beta am} \Gamma(0, m + \frac{a}{\beta}) - \Gamma(0, m) \right|$$

and since the absolute value expression is even in n

$$B = \sum_m \sum_{n=1}^{\infty} e^{-\frac{\pi\beta^2 n^2}{2\alpha}} e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} \left| e(-\frac{1}{2}n\beta a) e^{-\frac{1}{2}\pi\alpha a^2} e^{-\pi\alpha\beta am} \Gamma(n, m + \frac{a}{\beta}) - \Gamma(n, m) \right|.$$

Using (2.7) we can estimate B as follows

$$\begin{aligned} B &\leq \vartheta_3(0, 2i\alpha) \sum_m \sum_{n=1}^{\infty} e^{-\frac{\pi\beta^2 n^2}{2\alpha}} e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} (e^{-\frac{1}{2}\pi\alpha a^2} e^{-\pi\alpha\beta am} + 1) \\ &\quad \left[2\vartheta_3(i\frac{\pi}{2\alpha}\beta^2 n, it_\alpha) + e^{\pi\alpha/2} \vartheta_2(i\frac{\pi}{2\alpha}\beta^2 n, it_\alpha) \right] \\ &= \vartheta_3(0, 2i\alpha) M [2N_1 + e^{\pi\alpha/2} N_2] \end{aligned}$$

where

$$M = \sum_m e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} (e^{-\frac{1}{2}\pi\alpha a^2} e^{-\pi\alpha\beta am} + 1)$$

and

$$N_1 = \sum_{n=1}^{\infty} e^{-\frac{\pi\beta^2 n^2}{2\alpha}} \vartheta_3(i\frac{\pi}{2\alpha}\beta^2 n, it_\alpha), \quad N_2 = \sum_{n=1}^{\infty} e^{-\frac{\pi\beta^2 n^2}{2\alpha}} \vartheta_2(i\frac{\pi}{2\alpha}\beta^2 n, it_\alpha).$$

We will show that M is bounded over $\alpha > \frac{1}{4}$ and that $N_1 \rightarrow 0$ and $e^{\pi\alpha/2} N_2 \rightarrow 0$ as $q \rightarrow \infty$. For N_1 one uses Lemma A.2 to get (using $t_\alpha > 4$ for all $\alpha > 0$)

$$N_1 \leq (1 + t_\alpha^{-1/2}) \sum_{n=1}^{\infty} e^{-\pi \frac{\alpha\beta^2}{2(2\alpha^2+1)} n^2} < \frac{3}{2} \cdot \frac{1}{2} [\vartheta_3(0, \frac{i\alpha\beta^2}{2(2\alpha^2+1)}) - 1]$$

which clearly goes to 0 as $q \rightarrow \infty$ since $\frac{\alpha\beta^2}{2(2\alpha^2+1)} \rightarrow \infty$. For N_2 , one writes

$$N_2 = \sum_{n=1}^{\infty} e^{-\frac{\pi\beta^2 n^2}{2\alpha}} \sum_k e^{-\pi t_\alpha (k+\frac{1}{2})^2} e^{-\frac{\pi\beta^2 n}{\alpha} (k+\frac{1}{2})} = \sum_k e^{-\pi t_\alpha (k+\frac{1}{2})^2} R(k)$$

where, upon putting $c = \frac{\beta^2}{2\alpha}$, $d = k + \frac{1}{2}$, one has

$$\begin{aligned} R(k) &:= \sum_{n=1}^{\infty} e^{-\frac{\pi\beta^2 n^2}{2\alpha}} e^{-\frac{\pi\beta^2 n}{\alpha} (k+\frac{1}{2})} = \sum_{n=1}^{\infty} e^{-\pi c n^2} e^{-2\pi c d n} = e^{\pi c d^2} \sum_{n=1}^{\infty} e^{-\pi c (n+d)^2} \\ &< e^{\pi c d^2} \sum_{n=-\infty}^{\infty} e^{-\pi c (n+d)^2} = e^{\pi c d^2} \sum_{n=-\infty}^{\infty} e^{-\pi c (n+\frac{1}{2})^2} = e^{\pi c d^2} \vartheta_2(0, i\frac{\beta^2}{2\alpha}). \end{aligned}$$

Hence

$$N_2 < \sum_k e^{-\pi t_\alpha (k+\frac{1}{2})^2} e^{\pi \frac{\beta^2}{2\alpha} (k+\frac{1}{2})^2} \vartheta_2(0, i\frac{\beta^2}{2\alpha}) = \vartheta_2(0, 2i\alpha) \vartheta_2(0, i\frac{\beta^2}{2\alpha}).$$

Therefore, by (2.6)

$$e^{\pi\alpha/2} N_2 < e^{\pi\alpha/2} \vartheta_2(0, 2i\alpha) \vartheta_2(0, i\frac{\beta^2}{2\alpha}) < 2\vartheta_3(0, 2i\alpha) \vartheta_2(0, i\frac{\beta^2}{2\alpha})$$

which goes to zero (rapidly).

Next we show that M is bounded. Applying Lemma A.2,

$$\begin{aligned} M &= e^{-\frac{1}{2}\pi\alpha a^2} \sum_m e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} e^{-\pi\alpha m/q} + \sum_m e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} \\ &= e^{-\frac{1}{2}\pi\alpha a^2} \vartheta_3\left(\frac{i\pi\alpha}{2q}, i\frac{1}{2}\alpha\beta^2\right) + \vartheta_3(0, i\frac{1}{2}\alpha\beta^2) \\ &< \left[1 + \frac{\sqrt{2}}{\beta\sqrt{\alpha}}\right] e^{-\frac{1}{2}\pi\alpha a^2} \exp\left(\frac{\pi^2\alpha^2}{4q^2} \cdot \frac{2}{\pi\alpha\beta^2}\right) + \vartheta_3(0, i\frac{1}{2}\alpha\beta^2) \\ &= 1 + \frac{\sqrt{2}}{\beta\sqrt{\alpha}} + \vartheta_3(0, i\frac{1}{2}\alpha\beta^2) \end{aligned}$$

which converges to 2 as $q \rightarrow \infty$. Therefore, $B \rightarrow 0$. Next, we show that $A \rightarrow 0$. We have $A = |A_1| + A_2$ where

$$A_1 = e^{-\frac{1}{2}\pi\alpha a^2} \Gamma(0, \frac{a}{\beta}) - \Gamma(0, 0), \quad A_2 = \sum_{m \neq 0} e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} \left| e^{-\frac{1}{2}\pi\alpha a^2} e^{-\pi\alpha\beta a m} \Gamma(0, m + \frac{a}{\beta}) - \Gamma(0, m) \right|.$$

Using (2.6) and (2.7) again gives

$$\begin{aligned} A_2 &< \vartheta_3(0, 2i\alpha) [2\vartheta_3(0, it_\alpha) + e^{\pi\alpha/2} \vartheta_2(0, it_\alpha)] \sum_{m \neq 0} e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} \left[e^{-\pi\alpha m/q} e^{-\frac{1}{2}\pi\alpha a^2} + 1 \right] \\ &< 4\vartheta_3(0, 2i\alpha) \vartheta_3(0, it_\alpha) \sum_{m \neq 0} e^{-\frac{1}{2}\pi\alpha\beta^2 m^2} \left[e^{-\pi\alpha m/q} e^{-\frac{1}{2}\pi\alpha a^2} + 1 \right]. \end{aligned}$$

Expressing the summation as a ϑ_3 function and using Lemma A.2 one can show that it goes to zero as $q \rightarrow \infty$. Lastly, for A_1 , it will be enough to check that

$$\varepsilon := \kappa e^{\pi\alpha/2} \vartheta_2(t, 2i\alpha) - e^{\pi\alpha/2} \vartheta_2(0, 2i\alpha) \rightarrow 0$$

where $\kappa := e^{-\frac{1}{2}\pi\alpha a^2} < 1$ (which converges to 1 as $q \rightarrow \infty$) and $t = \frac{\pi}{2q}$. Since

$$e^{\pi\alpha/2} \vartheta_2(t, 2i\alpha) = e^{it} \sum_n e^{-2\pi\alpha n^2} e^{-2\pi\alpha n} e^{2itn}$$

one gets

$$\begin{aligned} \varepsilon &= \sum_n e^{-2\pi\alpha n^2} e^{-2\pi\alpha n} [\kappa e^{it} e^{2itn} - 1] \\ &= [\kappa e^{it} - 1] + [\kappa e^{-it} - 1] + \sum_{n \neq 0, -1} e^{-2\pi\alpha n^2} e^{-2\pi\alpha n} [\kappa e^{it} e^{2itn} - 1]. \end{aligned}$$

The first two terms here clearly go to zero, and the summation has absolute value no more than (since $n^2 + 2n \geq 0$ for all integers $n \neq -1$)

$$2 \sum_{n \neq 0, -1} e^{-2\pi\alpha n^2} e^{-2\pi\alpha n} = 2 \sum_{n \neq 0, -1} e^{-\pi\alpha n^2} e^{-\pi\alpha(n^2+2n)} < 2 \sum_{n \neq 0, -1} e^{-\pi\alpha n^2} < 2[\vartheta_3(0, i\alpha) - 1]$$

which clearly goes to zero. $\square$

We now have

$$\begin{aligned}
\|b^{-\frac{1}{2}}V_3^{p'}Xb^{-\frac{1}{2}} - V_3^{p'}\| &\leq \|b^{-\frac{1}{2}}V_3^{p'}Xb^{-\frac{1}{2}} - b^{-\frac{1}{2}}V_3^{p'}b^{\frac{1}{2}}\| + \|b^{-\frac{1}{2}}V_3^{p'}b^{\frac{1}{2}} - b^{-\frac{1}{2}}V_3^{p'}2^{\frac{1}{2}}\| \\
&\quad + \|b^{-\frac{1}{2}}V_3^{p'}2^{\frac{1}{2}} - 2^{-\frac{1}{2}}V_3^{p'}2^{\frac{1}{2}}\| \\
&\leq \|b^{-1}\|\|X - b\| + \|b^{-\frac{1}{2}}\|\|b^{\frac{1}{2}} - 2^{\frac{1}{2}}\| + 2^{\frac{1}{2}}\|b^{-\frac{1}{2}} - 2^{-\frac{1}{2}}\|.
\end{aligned}$$

Since the norms of b , b^{-1} , and $b^{\frac{1}{2}}$, are uniformly bounded by Lemma 3.2 (and Lemma A.3), and since $\|b - 2\| \rightarrow 0$, by Lemma 3.1, we conclude that $\|\langle \xi, U_1\xi \rangle_{D^\perp} - V_3^{p'}\| \rightarrow 0$ as $q \rightarrow \infty$. In the same fashion one shows that $\|\langle \xi, U_2\xi \rangle_{D^\perp} - V_4^{-p'}\| \rightarrow 0$, proving Proposition 4.1.

§5. Approximate Centrality

The main result of this section is the following, which proves (2) of the Main Theorem.

Theorem 5.1. *The Fourier orthogonal projection $e = \langle \xi, \xi \rangle_D$ (constructed in Section 3) is approximately central in A_θ . That is, for $j = 1, 2$, one has $\|U_j e - e U_j\| \rightarrow 0$ as $q \rightarrow \infty$.*

We begin by observing that the norm $\|U_1 e U_1^* - e\|$ can be made arbitrarily small provided that $\|U_1 \langle f, f \rangle_D U_1^* - \langle f, f \rangle_D\|$ can be made arbitrarily small. This follows from the basic fact that for any vector η in an equivalence A - B -bimodule and for $d, c \in B$ one has

$$\begin{aligned} \|\langle \eta d, \eta d \rangle_A - \langle \eta c, \eta c \rangle_A\| &\leq \|\langle \eta d, \eta d \rangle_A - \langle \eta d, \eta c \rangle_A\| + \|\langle \eta d, \eta c \rangle_A - \langle \eta c, \eta c \rangle_A\| \\ &= \|\langle \eta d, \eta(d - c) \rangle_A\| + \|\langle \eta(d - c), \eta c \rangle_A\| \\ &\leq \|\eta d\| \cdot \|\eta(d - c)\| + \|\eta(d - c)\| \cdot \|\eta c\| \\ &\leq \|\eta\|^2 (\|d\| + \|c\|) \|d - c\| \\ &= \|\langle \eta, \eta \rangle_B\| \cdot (\|d\| + \|c\|) \|d - c\| \end{aligned}$$

where $\|\eta\| = \|\langle \eta, \eta \rangle_B\|^{1/2} = \|\langle \eta, \eta \rangle_A\|^{1/2}$ (by Rieffel's Proposition 3.1 of [10]). Therefore, with $\eta = f, d = b^{-1/2}, c = 2^{-1/2}$ one has

$$\|e - \tfrac{1}{2} \langle f, f \rangle_D\| \leq \|b\| \left(\|b^{-1/2}\| + 2^{-1/2} \right) \|b^{-1/2} - 2^{-1/2}\|$$

which goes to zero as $q \rightarrow \infty$ by Lemmas 3.1 and 3.2. This also yields the eventual vanishing of $\|U_j e U_j^* - \tfrac{1}{2} U_j \langle f, f \rangle_D U_j^*\|$. Therefore, it is enough to show that

$$\|\langle f, f \rangle_D - U_j \langle f, f \rangle_D U_j^*\| \rightarrow 0$$

as $q \rightarrow \infty$ for $j = 1, 2$, so that the approximate centrality of the projection e follows.

One can check that

$$\langle f, f \rangle_D = \frac{\sqrt{2\alpha}}{\beta^2} \sum_{m,n} \overline{H(\frac{m}{\beta}, \frac{n}{\beta})} U_2^{qn} U_1^{qm}.$$

Hence

$$U_1 \langle f, f \rangle_D U_1^* - \langle f, f \rangle_D = \frac{\sqrt{2\alpha}}{\beta^2} \sum_{m,n} \overline{H(\frac{m}{\beta}, \frac{n}{\beta})} [\lambda^{qn} - 1] U_2^{qn} U_1^{qm}$$

and as $|H|$ is even in each variable one gets

$$\begin{aligned} \|U_1 \langle f, f \rangle_D U_1^* - \langle f, f \rangle_D\| &\leq \frac{\sqrt{2\alpha}}{\beta^2} \sum_{m,n} |H(\frac{m}{\beta}, \frac{n}{\beta})| |\lambda^{qn} - 1| \\ &= \frac{2\sqrt{2\alpha}}{\beta^2} \sum_m \sum_{n=1}^{\infty} |H(\frac{m}{\beta}, \frac{n}{\beta})| |\lambda^{qn} - 1| \end{aligned}$$

Since $\lambda^{qn} = e(qn\theta) = e(qn(\theta - \frac{p}{q})) = e(qna^2) = e(\frac{n}{q\beta^2})$ and since $|e(t) - 1| \leq 2\pi t$ for $t \geq 0$ one has $|\lambda^{qn} - 1| \leq \frac{2\pi n}{q\beta^2}$. Hence

$$\varepsilon_1 := \|U_1 \langle f, f \rangle_D U_1^* - \langle f, f \rangle_D\| \leq \frac{4\pi\sqrt{2\alpha}}{q\beta^4} \sum_m \sum_{n=1}^{\infty} n |H(\frac{m}{\beta}, \frac{n}{\beta})|.$$

Inserting

$$H\left(\frac{m}{\beta}, \frac{n}{\beta}\right) = \frac{1}{\sqrt{2\alpha}} e\left(-\frac{mn}{2\beta^2}\right) e^{-\frac{\pi\alpha}{2\beta^2}m^2} e^{-\frac{\pi}{2\alpha\beta^2}n^2} \Gamma\left(\frac{n}{\beta^2}, \frac{m}{\beta^2}\right)$$

and using the inequality (from (2.7))

$$|\Gamma\left(\frac{n}{\beta^2}, \frac{m}{\beta^2}\right)| \leq \vartheta_3(0, 2i\alpha) \left[2\vartheta_3\left(\frac{i\pi n}{2\alpha}, it_\alpha\right) + e^{\pi\alpha/2} \vartheta_2\left(\frac{i\pi n}{2\alpha}, it_\alpha\right) \right]$$

yields

$$\begin{aligned} \varepsilon_1 &\leq \frac{4\pi\vartheta_3(0, 2i\alpha)}{q\beta^4} \left[\sum_m e^{-\frac{\pi\alpha}{2\beta^2}m^2} \right] \sum_{n=1}^{\infty} n e^{-\frac{\pi}{2\alpha\beta^2}n^2} \left[2\vartheta_3\left(\frac{i\pi n}{2\alpha}, it_\alpha\right) + e^{\pi\alpha/2} \vartheta_2\left(\frac{i\pi n}{2\alpha}, it_\alpha\right) \right] \\ &= \frac{4\pi}{q\beta^4} \vartheta_3(0, 2i\alpha) \vartheta_3(0, i\frac{\alpha}{2\beta^2}) \sum_{n=1}^{\infty} n e^{-\frac{\pi}{2\alpha\beta^2}n^2} \left[2\vartheta_3\left(\frac{i\pi n}{2\alpha}, it_\alpha\right) + e^{\pi\alpha/2} \vartheta_2\left(\frac{i\pi n}{2\alpha}, it_\alpha\right) \right] \\ &= \frac{4\pi}{q\beta^4} \vartheta_3(0, 2i\alpha) \vartheta_3(0, i\frac{\alpha}{2\beta^2}) (2A_1 + B_1) \end{aligned}$$

where

$$A_1 = \sum_{n=1}^{\infty} n e^{-\frac{\pi}{2\alpha\beta^2}n^2} \vartheta_3\left(\frac{i\pi n}{2\alpha}, it_\alpha\right), \quad B_1 = e^{\pi\alpha/2} \sum_{n=1}^{\infty} n e^{-\frac{\pi}{2\alpha\beta^2}n^2} \vartheta_2\left(\frac{i\pi n}{2\alpha}, it_\alpha\right).$$

These are estimated by the following lemma, in the proof of which we make repeated implicit use of Lemmas A.1 and A.2 of the Appendix.

Lemma 5.2. *One has*

$$A_1 < \frac{9\beta^2}{2\pi\alpha} (2\alpha^2 + 1), \quad B_1 < \frac{12}{\pi} \alpha\beta^2 \vartheta_3(0, 2i\alpha) + 6\beta^2 (2 + \beta\sqrt{2\alpha}).$$

Proof. For A_1 , using Lemma A.2 one has

$$\vartheta_3\left(\frac{i\pi n}{2\alpha}, it_\alpha\right) < \frac{3}{2} \exp\left(\frac{\pi n^2}{4\alpha^2 t_\alpha}\right)$$

thus

$$A_1 < \frac{3}{2} \sum_{n=1}^{\infty} n e^{-\pi c n^2} < \frac{3}{2} \left(\frac{1}{2\pi c} + \frac{2}{\sqrt{2\pi e c}} \right)$$

where $c = -\frac{1}{2\alpha\beta^2} + \frac{1}{4\alpha^2 t_\alpha} = \frac{\alpha}{2\beta^2(2\alpha^2+1)}$. Since $2\pi c < 1$ (as $c < \frac{1}{8}$) and one obtains $\frac{1}{\sqrt{2\pi c}} < \frac{1}{2\pi c}$ one gets $A_1 < \frac{9}{4\pi c} = \frac{9\beta^2}{2\pi\alpha} (2\alpha^2 + 1)$.

To estimate B_1 a little more work is needed. One writes

$$B_1 = e^{\pi\alpha/2} \sum_{n=1}^{\infty} n e^{-\frac{\pi n^2}{2\alpha\beta^2}} \sum_k e^{-\pi t_\alpha (k+\frac{1}{2})^2} e^{-\frac{\pi n}{\alpha} (k+\frac{1}{2})} = e^{\pi\alpha/2} \sum_k e^{-\pi t_\alpha (k+\frac{1}{2})^2} N(k)$$

where

$$N(k) = \sum_{n=1}^{\infty} n e^{-\frac{\pi n^2}{2\alpha\beta^2}} e^{-\frac{\pi n}{\alpha}(k+\frac{1}{2})}.$$

To estimate $N(k)$, put $a = \frac{1}{2\alpha\beta^2}$ and $b = \beta^2(k + \frac{1}{2})$ to get

$$\begin{aligned} N(k) &= e^{\pi ab^2} \sum_{n=1}^{\infty} n e^{-\pi a(n+b)^2} \\ &< e^{\pi ab^2} \left[2|b| + \sqrt{\frac{2}{\pi a}} + \int_0^{\infty} x e^{-\pi a(x+b)^2} dx \right] \\ &< e^{\pi ab^2} \left[2|b| + \sqrt{\frac{2}{\pi a}} + \frac{1}{\pi a} + \frac{|b|}{\sqrt{a}} \right]. \end{aligned}$$

As $\alpha > \frac{1}{4}$ and $\beta > 2$, $a < \frac{1}{2}$, and thus $\sqrt{\frac{2}{\pi a}} < \frac{2}{\pi a}$ and we have

$$N(k) < e^{\frac{\pi\beta^2}{2\alpha}(k+\frac{1}{2})^2} \left[\frac{3}{\pi a} + |b| \left(2 + \frac{1}{\sqrt{a}} \right) \right].$$

Since $-t_\alpha + \frac{\beta^2}{2\alpha} = -2\alpha$, one gets

$$\begin{aligned} B_1 &< e^{\pi\alpha/2} \sum_k e^{-2\pi\alpha(k+\frac{1}{2})^2} \left[\frac{6}{\pi} \alpha\beta^2 + \beta^2(2 + \beta\sqrt{2\alpha})|k + \frac{1}{2}| \right] \\ &= \frac{6}{\pi} \alpha\beta^2 e^{\pi\alpha/2} \vartheta_2(0, 2i\alpha) + \beta^2(2 + \beta\sqrt{2\alpha})F(\alpha) \end{aligned}$$

where

$$\begin{aligned} F(\alpha) &= e^{\pi\alpha/2} \sum_k |k + \frac{1}{2}| e^{-2\pi\alpha(k+\frac{1}{2})^2} = 2e^{\pi\alpha/2} \sum_{k=0}^{\infty} (k + \frac{1}{2}) e^{-2\pi\alpha(k+\frac{1}{2})^2} \\ &= 2 \sum_{k=0}^{\infty} (k + \frac{1}{2}) e^{-2\pi\alpha(k^2+k)} \\ &< 2 \sum_{k=0}^{\infty} (k + \frac{1}{2}) e^{-2\pi\alpha k^2} = 1 + 2 \sum_{k=1}^{\infty} (k + \frac{1}{2}) e^{-2\pi\alpha k^2} \\ &< 1 + 4 \sum_{k=1}^{\infty} k e^{-2\pi\alpha k^2} < 1 + 4 \left[\frac{1}{4\pi\alpha} + \frac{1}{\sqrt{\pi e\alpha}} \right] < 6 \end{aligned}$$

since $\alpha > \frac{1}{4}$. Together with the inequality $e^{\pi\alpha/2} \vartheta_2(0, 2i\alpha) < 2\vartheta_3(0, 2i\alpha)$ one obtains the desired estimate on B_1 . $\square$

In view of this lemma, one now obtains

$$\varepsilon_1 < \frac{4\pi}{q\beta^2} \vartheta_3(0, 2i\alpha) \vartheta_3(0, i\frac{\alpha}{2\beta^2}) \left[\frac{9}{2\pi\alpha} (2\alpha^2 + 1) + \frac{12}{\pi} \alpha \vartheta_3(0, 2i\alpha) + 6(2 + \beta\sqrt{2\alpha}) \right]$$

and using the inequality $\vartheta_3(0, i\frac{\alpha}{2\beta^2}) \leq 1 + \frac{\beta\sqrt{2}}{\sqrt{\alpha}}$, it is now evident that all terms involving α and β are bounded over the interval $\alpha > \frac{1}{4}$, and therefore this upper bound vanishes as $q \rightarrow \infty$. To complete the proof of Theorem 5.1, we show (by similar ideas, but which we include for completeness) that $\varepsilon_2 = \|U_2 \langle f, f \rangle_D U_2^* - \langle f, f \rangle_D\|$ also vanishes as $q \rightarrow \infty$. We have

$$\begin{aligned} \varepsilon_2 &\leq \frac{\sqrt{2\alpha}}{\beta^2} \sum_{m,n} |H(\frac{m}{\beta}, \frac{n}{\beta})| |\lambda^{-qm} - 1| = \frac{2\sqrt{2\alpha}}{\beta^2} \sum_n \sum_{m=1}^{\infty} |H(\frac{m}{\beta}, \frac{n}{\beta})| |\lambda^{qm} - 1| \\ &= \frac{2}{\beta^2} \sum_n \sum_{m=1}^{\infty} e^{-\frac{\pi\alpha}{2\beta^2}m^2} e^{-\frac{\pi}{2\alpha\beta^2}n^2} |\Gamma(\frac{n}{\beta^2}, \frac{m}{\beta^2})| |\lambda^{qm} - 1| \\ &\leq \frac{4\pi}{q\beta^4} \sum_n \sum_{m=1}^{\infty} m e^{-\frac{\pi\alpha}{2\beta^2}m^2} e^{-\frac{\pi}{2\alpha\beta^2}n^2} |\Gamma(\frac{n}{\beta^2}, \frac{m}{\beta^2})| \\ &\leq \frac{4\pi}{q\beta^4} \vartheta_3(0, 2i\alpha) \left(\sum_{m=1}^{\infty} m e^{-\frac{\pi\alpha}{2\beta^2}m^2} \right) \sum_n e^{-\frac{\pi}{2\alpha\beta^2}n^2} \left[2\vartheta_3(\frac{i\pi n}{2\alpha}, it_\alpha) + e^{\pi\alpha/2} \vartheta_2(\frac{i\pi n}{2\alpha}, it_\alpha) \right] \end{aligned}$$

As before, let

$$A_2 = \sum_n e^{-\frac{\pi}{2\alpha\beta^2}n^2} \vartheta_3(\frac{i\pi n}{2\alpha}, it_\alpha), \quad B_2 = e^{\pi\alpha/2} \sum_n e^{-\frac{\pi}{2\alpha\beta^2}n^2} \vartheta_2(\frac{i\pi n}{2\alpha}, it_\alpha).$$

First note that by Lemma A.1,

$$\frac{\alpha}{\beta^2} \sum_{m=1}^{\infty} m e^{-\frac{\pi\alpha}{2\beta^2}m^2} < \frac{\alpha}{\beta^2} \left[\frac{\beta^2}{\pi\alpha} + \frac{2\beta}{\sqrt{\pi e\alpha}} \right] < \frac{1}{\pi} + \frac{\sqrt{\alpha}}{\beta} < 2$$

for $\alpha > \frac{1}{4}$. Thus, $\varepsilon_2 \rightarrow 0$ follows once it is shown that $\frac{1}{\alpha\beta^2}(2A_2 + B_2)$ is bounded for $\alpha > \frac{1}{4}$ (and $\vartheta_3(0, 2i\alpha)$ being already bounded over this range). Since $t_\alpha = 4\alpha + \frac{2}{\alpha} > 4$ for $\alpha > 0$,

$$A_2 = \vartheta_3(0, it_\alpha) + 2 \sum_{n=1}^{\infty} e^{-\frac{\pi}{2\alpha\beta^2}n^2} \vartheta_3(\frac{i\pi n}{2\alpha}, it_\alpha) < \vartheta_3(0, 4i) + 2A_1 < \vartheta_3(0, 4i) + \frac{9\beta^2}{\pi\alpha}(2\alpha^2 + 1)$$

(by Lemma 5.2), thus clearly $\frac{A_2}{\alpha\beta^2}$ is bounded. Finally,

$$B_2 = e^{\pi\alpha/2} \vartheta_2(0, it_\alpha) + 2e^{\pi\alpha/2} \sum_{n=1}^{\infty} e^{-\frac{\pi}{2\alpha\beta^2}n^2} \vartheta_2(\frac{i\pi n}{2\alpha}, it_\alpha) \quad (*)$$

and upon inserting the summation for ϑ_2 as we have done above for $N(k)$ (which this time does not include the 'n' in front of the exponential), one gets that the summation in (*) is approximately no more than $\beta\sqrt{2\alpha}\vartheta_2(0, 2i\alpha)$. Hence the second term in (*) is no more than $2\beta\sqrt{2\alpha}e^{\pi\alpha/2}\vartheta_2(0, 2i\alpha) < 4\beta\sqrt{2\alpha}\vartheta_3(0, 2i\alpha)$, which now makes it clear that $\frac{B_2}{\alpha\beta^2}$ is bounded (in fact goes to zero) over $\alpha > \frac{1}{4}$. This proves that $\varepsilon_2 \rightarrow 0$ as $q \rightarrow \infty$ and completes the proof of Theorem 5.1.

§6. Appendix

The following lemmas are easy but are included for completeness.

Lemma A.1. *For all $a > 0$ and b real, one has*

$$\begin{aligned} \sum_{n=1}^{\infty} n e^{-\pi a(n+b)^2} &< \sqrt{\frac{2}{\pi a}} + \frac{1}{\pi a} + 2|b| + \frac{|b|}{\sqrt{a}} \\ \sum_{n=1}^{\infty} n e^{-\pi a n^2} &< \frac{1}{2\pi a} + \frac{2}{\sqrt{2\pi e a}}. \end{aligned}$$

Proof. Let $f(x) = x e^{-\pi a(x+b)^2}$. Then f is increasing on $[0, x_0]$ and decreasing on $[x_0, \infty)$, where $x_0 = \frac{1}{2}(\sqrt{b^2 + \frac{2}{\pi a}} - b)$. Let $N \geq 0$ be the greatest integer $\leq x_0$. Then

$$\sum_{n=1}^{N-1} f(n) < \int_0^N f(x) dx, \quad \sum_{n=N+2}^{\infty} f(n) < \int_{N+1}^{\infty} f(x) dx.$$

Hence

$$\sum_{n=1}^{\infty} f(n) < \int_0^N f(x) dx + \int_{N+1}^{\infty} f(x) dx + f(N) + f(N+1) < 2f(x_0) + \int_0^{\infty} f(x) dx.$$

The first inequality holds from $2f(x_0) < 2x_0 < 2|b| + \sqrt{\frac{2}{\pi a}}$ and

$$\begin{aligned} \int_0^{\infty} x e^{-\pi a(x+b)^2} dx &< \int_{-\infty}^{\infty} x e^{-\pi a(x+b)^2} dx = \int_{-\infty}^{\infty} |x-b| e^{-\pi a x^2} dx \\ &\leq \int_{-\infty}^{\infty} |x| e^{-\pi a x^2} dx + |b| \int_{-\infty}^{\infty} e^{-\pi a x^2} dx \\ &= 2 \int_0^{\infty} x e^{-\pi a x^2} dx + \frac{|b|}{\sqrt{a}} = \frac{1}{\pi a} + \frac{|b|}{\sqrt{a}}. \end{aligned}$$

Similarly, for the second equality note that the function $x e^{-\pi a x^2}$ has the maximum value $\frac{1}{\sqrt{2\pi e a}}$ so that

$$\sum_{n=1}^{\infty} n e^{-\pi a n^2} < \frac{2}{\sqrt{2\pi e a}} + \int_0^{\infty} x e^{-\pi a x^2} dx = \frac{2}{\sqrt{2\pi e a}} + \frac{1}{2\pi a}.$$

□

Lemma A.2. *For $A > 0$ and any real number B one has*

$$0 < \vartheta(iB, iA) \leq \left(1 + \frac{1}{\sqrt{A}}\right) e^{B^2/(\pi A)}$$

for $\vartheta = \vartheta_2, \vartheta_3$.

Proof. Let $C = B/(\pi A)$, let N be the greatest integer $\leq C$, and put

$$\xi = \min(C - N, N + 1 - C) \leq \frac{1}{2}.$$

From a graphical consideration of areas one has

$$\sum_p e^{-\pi A(p-C)^2} \leq e^{-\pi A\xi^2} + \int_{\mathbb{R}} e^{-\pi A(x-C)^2} dx \leq 1 + \frac{1}{\sqrt{A}}$$

which holds for any real C . Hence,

$$\vartheta_3(iB, iA) = \sum_p e^{-\pi A p^2} e^{2Bp} = e^{B^2/(\pi A)} \sum_p e^{-\pi A(p-C)^2} \leq \left(1 + \frac{1}{\sqrt{A}}\right) e^{B^2/(\pi A)}$$

A similar argument holds for ϑ_2 , hence the result. $\square$

Lemma A.3. *If x and y are positive invertible elements in a unital C^* -algebra, then*

$$\|x^{\frac{1}{2}} - y^{\frac{1}{2}}\| \leq \sqrt{2} m^{-2} M^{\frac{3}{2}} \|x - y\|$$

where $m = \min(\|x^{-1}\|^{-1}, \|y^{-1}\|^{-1})$ and $M = \max(\|x\|, \|y\|)$.

Proof. Since the smallest and largest points of the spectrum of x are $\|x^{-1}\|^{-1}$ and $\|x\|$, respectively, the spectra of x and y are contained in the interval $[m, M]$. Let ϵ, c be arbitrary positive numbers with $\epsilon < m$ and $M < c$. If γ is the circle centered at c of radius $c - \epsilon$, then it is contained in the right half of the complex plane and surrounds $[m, M]$. Taking the principal branch of the square root function, one has

$$x^{\frac{1}{2}} - y^{\frac{1}{2}} = \frac{1}{2\pi i} \int_{\gamma} z^{\frac{1}{2}} [(z - x)^{-1} - (z - y)^{-1}] dz = \frac{1}{2\pi i} \int_{\gamma} z^{\frac{1}{2}} (z - x)^{-1} (x - y) (z - y)^{-1} dz$$

and since $|z^{\frac{1}{2}}| \leq (2c - \epsilon)^{\frac{1}{2}}$ and $\|(z - x)^{-1}\| \leq (m - \epsilon)^{-1}$ (and also for y) for all z on γ and $\int_{\gamma} |dz| = 2\pi(c - \epsilon)$, one obtains

$$\|x^{\frac{1}{2}} - y^{\frac{1}{2}}\| \leq (m - \epsilon)^{-2} (c - \epsilon) (2c - \epsilon)^{\frac{1}{2}} \|x - y\|.$$

Letting $\epsilon \rightarrow 0$ and $c \rightarrow M$ yields the result. $\square$

References

- [1] F. P. Boca, *On the flip fixed point algebra in certain noncommutative tori*, Indiana J. Math. **45** (1996), 253–272.
- [2] ———, *Projections in rotation algebras and theta functions*, Comm. Math. Phys. **202** (1999), 325–357.
- [3] O. Bratteli, G. A. Elliott, D. E. Evans, A. Kishimoto, *Non-commutative spheres I*, Internat. J. Math. **2** (1990), no. 2, 139–166.
- [4] O. Bratteli, G. A. Elliott, D. E. Evans, A. Kishimoto, *Non-commutative spheres II: rational rotations*, J. Operator Theory **27** (1992), 53–85.
- [5] O. Bratteli and A. Kishimoto, *Non-commutative spheres III. Irrational Rotations*, Comm. Math. Phys. **147** (1992), 605–624.
- [6] G. Elliott and D. Evans, *The structure of the irrational rotation C^* -algebra*, Ann. Math. **138** (1993), 477–501.
- [7] G. Elliott and Q. Lin, *Cut-down method in the inductive limit decomposition of non-commutative tori*, J. London Math. Soc. (2) **54** (1996), no. 1, 121–134.
- [8] A. Kumjian, *On the K -theory of the symmetrized non-commutative torus*, C. R. Math. Rep. Acad. Sci. Canada **12** (1990), no. 3, 87–89.
- [9] S. Lang, *Fundamentals of Diophantine geometry*, Springer-Verlag, New York-Berlin, 1983.
- [10] M. Rieffel, *Unitary representations of group extensions; an algebraic approach to the theory of Mackey and Blattner*, Studies in Analysis, Adv. Math. Suppl. Series **4** (1979), 43–82.
- [11] ———, *C^* -algebras associated with irrational rotations*, Pacific J. Math. **93** (1981), no. 2, 415–429.
- [12] ———, *Projective modules over higher-dimensional non-commutative tori*, Canad. J. Math. **40** (1988), 257–338.
- [13] K. F. Roth, *Rational approximations to algebraic numbers*, Mathematika **2** (1955), 1–20; corrigendum, 168.
- [14] S. G. Walters, *Inductive limit automorphisms of the irrational rotation algebra*, Comm. Math. Phys. **171** (1995), 365–381.
- [15] ———, *Chern characters of Fourier modules*, Canad. J. Math. **52** (2000), no. 3, 633–672.
- [16] ———, *K -theory of non commutative spheres arising from the Fourier automorphism*, Canad. J. Math. **53** (2001), no. 3, 631–672.
- [17] ———, *On the inductive limit structure of order four automorphisms of the irrational rotation algebra*, Internat. J. Math. (2001), 8 pages (to appear).
- [18] ———, *On Fourier orthogonal projections in the rotation algebra*, preprint (2000), 16 pages.
- [19] E. T. Whittaker and G. N. Watson, *A course in modern analysis*, Cambridge Univ. Press, 1950 (4th edition).

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, THE UNIVERSITY OF NORTHERN BRITISH COLUMBIA, PRINCE GEORGE, B.C. V2N 4Z9 CANADA

E-mail address: `walters@hilbert.unbc.ca` or `walters@unbc.ca`

Website: `http://hilbert.unbc.ca/walters`